

**CM20019—COMPUTATION III:
FORMAL LOGIC AND SEMANTICS
COURSEWORK, DUE 22.11.2007**

ALESSIO GUGLIELMI

Information and instructions for the coursework submission process:

- This coursework is to be submitted through the coursework post box located outside 1W2.23 by 22 November 2007 at 17:00. Please be careful to select the box relative to this course.
- This coursework is to be submitted in written form, on paper, bound securely. Please make sure it contains information to identify you, clearly written: surname, first name, and student number. You must include the coursework signature sheet.
- This coursework will contribute 25% of the final grade for this unit.
- This is an individual coursework, and it should be completed within the student's own time.
- Feedback will be provided by example solutions.

Plagiarism will be detected and will be penalised. Remember that, while there are not many ways to answer a problem correctly, there are infinitely many, **unique** ways of making mistakes. Remember also that **even the best make mistakes**.

* * *

Problem 1. Use the unification algorithm on the following pairs of terms and either find most general unifiers or tell when the terms are not unifiable and why.

- (a) $f(X, Y)$ and $f(Z, X)$; [2 points]
- (b) $f(X, Y)$ and $g(Z, X)$; [2 points]
- (c) $f(X, X)$ and $f(Y, g(Y))$; [3 points]
- (d) $f(X, Y, g(g(Z)))$ and $f(g(Y), g(Z), X)$; [4 points]
- (e) $f(X, g(a))$ and $f(Y, X)$. [4 points]

Problem 2. Either prove the validity, by resorting to relevant definitions, or disprove it, by providing a counter-model, of the following formulae:

- (a) $F = (\forall X)(\exists Y)(p(X) \rightarrow p(Y))$; [6 points]
- (b) $G = (\exists X)(\forall Y)(p(X) \rightarrow p(Y))$. [9 points]

Then, answer the question:

- (c) Are F and G equivalent? [5 points]

(Note: a counter-model for a formula is an interpretation in which the formula is false; two formulae are equivalent if whenever one is true in an interpretation, the other is true in that interpretation as well.)

Problem 3. We are given the predicate symbols p and q , and the function symbol f , all of which have arity 1. Let I be an interpretation such that:

- its domain is the set \mathbb{N} of natural numbers (including 0);
- p is interpreted as $\{3n \mid n \in \mathbb{N}\} \subset \mathbb{N}$;
- q is interpreted as $\{n \mid n \text{ is prime}\} \subset \mathbb{N}$;
- f is interpreted as the function $\mathbf{f}: \mathbb{N} \rightarrow \mathbb{N}$ such that $\mathbf{f}(n) = n + 6$.

Please answer the following questions in concise, not necessarily formal, but precise and unambiguous mathematical terms.

- (a) Is the formula $p(f(X)) \wedge q(X)$ false in I for some valuation? If so, which one? If not, why not? [2 points]
- (b) Is the formula $p(f(X)) \wedge q(X)$ true in I for some valuation? If so, which one? If not, why not? [2 points]
- (c) Is the formula $\neg(\exists X)(p(f(X)) \wedge q(X))$ true in I ? Explain. [2 points]
- (d) Is the formula $(\forall X)(q(X) \rightarrow (p(f(X)) \vee q(f(X))))$ true in I ? Explain. [5 points]
- (e) Is the formula $(\forall X)(q(X) \rightarrow (p(f(X)) \vee q(f(X))))$ valid? Explain. [4 points]
- (f) Find, in the given language, a satisfiable formula F such that it contains the symbol f and such that I is a model for

$$(\forall X)(F \rightarrow (p(X) \wedge q(X))) \quad .$$

[11 points]

- (g) Find, in the given language, a formula F such that it contains the symbol f and such that

$$(\forall X)(F \rightarrow (p(X) \wedge q(X)))$$

is valid.

[9 points]

(Note: a formula F is satisfiable if there is an interpretation and a valuation for which F is true; the set of prime numbers is $\{2, 3, 5, 7, 11, 13, 17, \dots\}$.)

Problem 4. Consider the following Prolog program:

$a(0, X, X)$.

$a(s(X), Y, s(Z)) \text{ :- } a(X, Y, Z)$.

$b(0, 0)$.

$b(s(0), s(0))$.

$b(s(s(0)), s(0))$.

$b(s(s(s(Y))), X) \text{ :- } b(Y, Z),$
 $\quad b(s(Y), W),$
 $\quad b(s(s(Y)), U),$
 $\quad a(Z, W, V),$
 $\quad a(V, U, X)$.

$c(X) \text{ :- } b(Y, X)$.

$d(X) \text{ :- } b(X, X)$.

In your answers, please abbreviate any string of the kind $s(\dots s(0)\dots)$ with the numeral corresponding to the number of its s symbols; for example, please abbreviate $s(s(s(0)))$ as 3. Please explain the meaning of predicates in concise, not necessarily formal, but precise and unambiguous mathematical terms.

- (a) What is the meaning of the predicate $a(X, Y, Z)$? [6 points]
- (b) What is the meaning of the predicate $b(X, Y)$? [16 points]
- (c) What is the meaning of the predicate $c(X)$? [4 points]
- (d) What is the meaning of the predicate $d(X)$? [4 points]